



Group Meeting Problems 04/13/2009


Solution: Kikuchi





Langer, P. Chem. Commun. 2003, 3094







Can. J. Chem. 79, 2001, 1694 / org. lett 2003, 4/63




Chem. Commun 1909 (1996)
G.W. Gribble



5





Tetrahedron Lett. Vol 33, p6849 (1992) M. Terashima




If more than 2 eq. KHMDS was used, the following mechanism would be suggested:




Stuart L. Schreiber, J. Am. Chem. Soc., 1989, 111, 2331




Groot A., et al., J. Org. Chem., 2001, 66, 2350




3. (Continued)




Org. Lect., 2002, 5, 3157



W. Dauben, J. Am. Chem. Soc., 1973, 95, 5088




Perkin Trans I, 2001 (23), 3150





Group Meeting Problems

Date 2004. 4. 21 No. 1



Katritzky, A.R. JCS. PT1. 1976, 2334







H^+ P. \sim




JOC (1992), 3085, M. Malacria.

3.




TL. 1973, 26, 2351 Y. Kita

4.




Group Meeting Problems

Date 2004. 4. 21 No. 2



S.M. Weinreb et al

Org. Lett. 3, 3507 (2001)



TCC, 1988, 53, 1116, Weinreb et al